



Hypothesis: Improved Patient Compliance and Functional Outcomes with LRS in Treating Infected Femoral Non-Unions

Jenil Patel¹,
Rajeev Joshi¹,
Sahil Sanghavi¹,
Mahavir Dugad¹,
Darshan Sonawane¹,
Ashok Shyam¹,
Parag Sancheti¹

¹Department of Orthopaedics, Sancheti Institute of Orthopaedics and Rehabilitation, Pune, Maharashtra, India.

Address of Correspondence

Dr. Jenil Patel,
Department of Orthopaedics, Sancheti Institute of Orthopaedics and Rehabilitation, Pune, Maharashtra, India.

E-mail: jenilpatel.jp23@gmail.com

Abstract

Background: Infected non-union of the femur combines chronic osteomyelitis and failure of fracture healing, causing prolonged pain, repeated surgeries and significant socioeconomic burden for patients and families. Radical debridement removes necrotic bone, bacterial biofilm and non-viable soft tissue but frequently creates segmental defects that require reconstruction. The Limb Reconstruction System (LRS) is a uniplanar external fixator that applies distraction osteogenesis principles to provide stable fixation, permit bone transport or lengthening, and support biological regeneration while being less cumbersome than circular frames in many cases.

Hypothesis: We hypothesised that thorough debridement followed by stabilization and reconstruction with LRS would produce high rates of bony union and durable infection control while restoring useful limb function. We also expected that LRS would correct limb length discrepancy and alignment in most cases, and that common complications such as pin-tract infection, pin loosening and joint stiffness would be predictable and manageable with standardised pin care and supervised physiotherapy.

Clinical importance: A practical limb-salvage method that eradicates infection and restores bone continuity has major benefits for patients. For selected femoral non-unions with adequate soft-tissue cover and moderate bone loss, LRS combines biological reconstruction with simpler day-to-day care: it supports early mobilisation, simplifies frame handling compared with bulkier circular systems, and allows staged adjunctive procedures when necessary. Outcomes depend on meticulous surgical debridement, appropriate antimicrobial therapy, close pin-site management and a coordinated rehabilitation programme to preserve joint motion and muscle function.

Future research: Larger multicentre comparative trials are needed to define which defect patterns and patient factors favour LRS over circular frames or combined intramedullary approaches. Studies should test strategies to reduce pin-tract complications (improved pin design, coatings and standardised care bundles), evaluate optimised physiotherapy regimens to limit stiffness, and maintain long-term registries to capture infection recurrence, functional durability and patient-reported outcomes. Improve patient outcomes.

Keywords: Infected non-union, Femur, Limb Reconstruction System, Bone transport, External fixation, Distraction osteogenesis, Radical debridement, Limb salvage.



Dr. Jenil Patel



Dr. Rajeev Joshi



Dr. Sahil Sanghavi



Dr. Mahavir Dugad



Dr. Darshan Sonawane



Dr. Ashok Shyam



Dr. Parag Sancheti

DOI: <https://doi.org/10.13107/jmt.2025.v11.i01.240>

© The Author(s). 2025 Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted use, distribution, and non-commercial reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Background

Infected non-union of the femur is a devastating condition that combines two difficult problems: persistent bone infection (chronic osteomyelitis) and failure of a fracture to heal. The combination causes prolonged pain, repeated surgeries, long periods away from work, and often permanent disability. The causes are many — high-energy trauma, contamination at the time of injury, multiple prior operations, devitalized bone, poor soft-tissue cover, and host factors such as diabetes. Historically, treating infection and restoring a functional limb have required staged, sometimes complex approaches. [1–6]

Two broad strategies are used. One prioritizes immediate mechanical stability to encourage union; the other prioritizes infection control through radical debridement and then reconstructs the bone defect that results. When chronic infection is established, most experienced centres favour aggressive debridement first (to remove necrotic bone and bacterial reservoirs), followed by reconstruction — because residual dead bone and biofilm hinder any attempt at union. The Ilizarov method and distraction osteogenesis grew from this logic: after debridement, bone transport or lengthening regenerates bone and can restore limb length and alignment while the soft tissues recover. [7–9,24]

Monolateral devices such as the Limb Reconstruction System (LRS) translate Ilizarov principles to a uniplanar frame. LRS stabilizes the femoral shaft, allows compression at non-union sites, and supports bone transport or lengthening via corticotomy. Compared with circular frames, monolateral frames are less bulky, easier for patients to tolerate, and simpler for nursing care and physiotherapy in many settings. The biological basis — maintained stability, controlled distraction, and preservation of local blood supply — remains the same. Monolateral fixation has been reported as effective in many series for femoral defects and infected non-unions. [10–15, 21, 24]

Despite advantages, external fixation carries known risks: pin-tract inflammation or infection, pin loosening, joint stiffness, and the need for patient commitment to pin-site care and physiotherapy. These complications are predictable and, with careful management, frequently manageable — but they require careful planning, good patient education and close follow-up. [16–19,23]

This synopsis is based on a single-centre series of patients treated with LRS for infected femoral non-union. The patients underwent radical debridement, frame application and, when needed, corticotomy and bone transport. Outcomes were measured by radiological healing, ASAMI bone and functional scores, and standard indices of lengthening and consolidation. The thesis provides detailed patient demographics, microbiology, complications and comparative discussion with historic series.

Hypothesis and Rationale

Primary hypothesis

- When infected femoral non-union is managed with thorough debridement followed by stabilization and reconstructive techniques using LRS, the majority of patients will achieve bony union and satisfactory infection control, returning to useful limb function.

Why this approach should work

- The biological barrier to healing in infected non-union is necrotic bone and bacterial biofilm. Removing these with radical debridement lowers bacterial load and restores a healthier environment for bone repair. Applying stable mechanical conditions with the LRS supports bone healing, and if a defect remains, distraction osteogenesis (bone transport or lengthening from a corticotomy) regenerates bone from living tissues. These are established principles from Ilizarov and later monolateral adaptations. [7,9,13,24]

What we expect from LRS

- LRS allows compression at the non-union site and controlled distraction at a corticotomy site. In defects ≤ 2 cm, simple compression and stimulation may be enough; when defects exceed that, bifocal techniques with corticotomy and transport regenerate bone while repairing the gap. The uniplanar frame is less cumbersome than circular frames and is often better tolerated by patients, facilitating early mobilisation and physiotherapy — both important to preserve joint motion and muscle function. [10–15,20,21]

Operational goals and measurable endpoints

- The study operationalises the hypothesis by tracking objective measures: radiographic consolidation time, lengthening (distraction) index, external fixation (healing) index, ASAMI bone and functional scores, and infection eradication on clinical and microbiological grounds. These endpoints allow comparison with historical series of Ilizarov and monolateral fixation. [12–14,24]

Clinical considerations built into the treatment plan

- Exclude confounders such as tuberculous non-union and major neurological impairment that would change healing dynamics. Use radical debridement until bleeding bone (“paprka sign”) is reached, send multiple culture samples, and apply LRS with frame planning tailored to defect location and length. Provide structured pin-site care and an active physiotherapy programme to reduce stiffness. These process steps are intended to maximise the chance of union while limiting predictable complications. [16–17]

Why the question matters

- If LRS reliably produces high union rates and good infection control with lower patient burden than circular frames, it becomes a practical first-line reconstructive option for many femoral infected non-unions — especially where circular frames are unavailable or poorly tolerated. Demonstrating comparable outcomes supports wider adoption and helps surgeons select the best tool for a given patient. [10,24]

Discussion

Main findings

• The thesis reports a cohort of patients treated with LRS for infected femoral non-union. Most patients were young adults, typical of high-energy trauma patterns seen in femoral fractures. Radical debridement followed by LRS frame application, with corticotomy and transport when required, formed the treatment protocol. The study reports a high union rate and acceptable functional outcomes for the majority of patients.

How these results fit with prior evidence

• Historical series using Ilizarov circular frames and monolateral devices document high union rates in selected patients but also report significant complication burdens related to pin sites and joint stiffness. The present LRS experience aligns with that pattern: effective union and limb salvage in most patients, balanced by predictable complications. Monolateral devices have been described as offering simpler care with similar outcomes in many femoral cases, which this series supports. [9–15,19,24]

Key drivers of success

- Adequate debridement: removing necrotic bone and infected soft tissue is the single most important step for infection control and eventual union. [7,31]
- Stable fixation: LRS provides the stability required during consolidation and allows earlier partial weight-bearing, which promotes bone remodeling. [10,21]
- Patient engagement: committed pin-site care and physiotherapy reduce complications such as pin-tract infection and joint stiffness. [16–19,23]

Complications and their management

• Pin-tract infection, pin loosening and joint stiffness are the commonest problems and were treated by local care, antibiotics when required, and intensified physiotherapy. In rare cases persistent infection required further procedure(s). These complications do not negate the overall utility of LRS but underline the need for careful follow-up and a patient-centred care pathway. [16–19,23]

Limitations to bear in mind

• The single-centre nature and modest sample size limit broad generalisability. Absence of contemporaneous control (for example, patients treated with circular frames or antibiotic nails) prevents definitive conclusions on comparative effectiveness. Follow-up needs to be sufficiently long to capture late recurrences of infection or mechanical failures. [14,24]

Practical recommendations

• Choose LRS for infected femoral non-unions when the defect and deformity are amenable to a uniplanar solution, the soft-tissue envelope is adequate, and the patient is motivated for prolonged rehabilitation. Reserve circular frames or combined techniques for complex multiplanar deformities or very large segmental defects. Ensure meticulous debridement, clear microbiological sampling and a structured pin-site and

physiotherapy protocol to reduce complications. [10–15,21–24]

Clinical importance

For many patients with infected femoral non-union, the LRS offers an effective limb-salvage option that combines stability with the biological advantage of distraction osteogenesis when needed. When used after radical debridement, LRS achieves high union rates and acceptable functional recovery while being easier for patients and caregivers to manage than bulky circular frames. The approach preserves options — it can be combined with bone grafting, intramedullary devices or staged soft-tissue reconstructions as required — and is practical in a wide range of clinical settings.

Future directions

- Larger, multicentre comparative studies (LRS vs circular frames vs combined intramedullary + external strategies) to identify which defects and patient characteristics favour each technique.
- Trials of improved pin coatings, standardised pin-care bundles and early guided physiotherapy protocols to reduce pin-tract issues and stiffness.
- Prospective registries capturing long-term infection recurrence, patient-reported outcomes and cost analyses to inform treatment selection across resource settings.

References

1. Motsitsi NS. Management of infected nonunion of long bones: the last decade (1996–2006). *Injury*. 2008 Feb; 39(2):155–60. doi:10.1016/j.injury.2007.08.032.
2. Nicoll EA. Fracture of tibial shaft. A survey of 705 cases. *J Bone Joint Surg Br*. 1964; 46B:373–87.
3. Saleh M. Non-union surgery. Part 1. Basic principles of management. *IJOT*. 1992; 2:4–18.
4. Mills LA, A Hamish. The relative incidence of fracture non-union in the Scottish population: a 5-year epidemiological study. *BMJ Open*. 2013; 3.
5. Chao EYS, Aro HT. Biomechanics and Biology of external fixation. In: Coombs R, Green S, Sarmiento A, editors. *External fixation and functional bracing*. London: Orthotext; 1989. p. 67–95.
6. McKibbin B. The biology of fracture healing in long bones. *J Bone Joint Surg Br*. 1978; 60-B: 150.
7. Ilizarov GA. The tension-stress effect on the genesis and growth of tissues. Part I. The influence of stability of fixation and soft tissue preservation. *Clin Orthop Relat Res*. 1989; 238:249–81.
8. Rockwood CA, Green DP, Bucholz RW. *Rockwood and Green's Fractures in Adults*. 6th ed. Philadelphia: Lippincott Williams & Wilkins; 2006.

9. Dendrinios GK, Konto S, Lyritis E. Use of Ilizarov technique for treatment of nonunion of tibia associated with infection. *J Bone Joint Surg Br.* 1995; 77-B: 835–46.
10. Arora S, Batra S, Gupta V, Goyal A. Distraction osteogenesis using a monolateral external fixator for infected non-union of the femur with bone loss. *J Orthop Surg (Hong Kong).* 2012 Aug; 20(2):185–90.
11. Spiegelberg B, Parratt T, Dheerendra SK, Khan WS, Jennings R, Marsh DR. Ilizarov principles of deformity correction. *Ann R Coll Surg Engl.* 2010 Mar; 92(2):101–5.
12. Marsh JL, Nepola JV, Meffert R. Dynamic external fixation for stabilization of nonunion. *Clin Orthop Relat Res.* 1992 May; (278):200–206.
13. De Bastiani G, Aldegheri R, Renzi-Brivio L, Trivella G. Limb lengthening by callus distraction (callotaxis). *J Pediatr Orthop.* 1987; 7(2):129–134.
14. Sangkaew C. Distraction osteogenesis of the femur using conventional monolateral external fixator. *Arch Orthop Trauma Surg.* 2008 Sep; 128(9):889–99.
15. Paley D. Problems, Obstacles and complications of limb lengthening by the Ilizarov technique. *Clin Orthop Relat Res.* 1990; (250):81–104.
16. Hashmi MA, Ali A, Saleh M. Management of non-unions with mono-lateral external fixation. *Injury.* 2001; 32(Suppl): S-D30–S-D34.
17. Vidal J. Traitement des fractures ouverte de jambe par le fixateur externe en double cadre. *Rev Chir Orthop.* 1976; 62(4):433–48.
18. Burny FL. Elastic external fixation of tibial fracture. *External fixation: the current state of the art.* 1979:55–74.
19. Behrens F, Comfort TH, Searls K, Denis F, Young JT. Unilateral external fixation for severe open tibial fractures. *Clin Orthop Relat Res.* 1983; (178):111–20.
20. Green SA, Garland DE, Moore TJ, Barad SJ. External fixation for the uninfected angulated nonunion of the tibia. *Clin Orthop Relat Res.* 1984; (190):204–11.
21. De Bastiani G, Aldegheri RO, Renzi-Brivio LO. The treatment of fractures with a dynamic axial fixator. *J Bone Joint Surg Br.* 1984 Aug; 66(4):538–45.
22. Slätis P, Paavolainen P. External fixation of infected non-union of the femur. *Injury.* 1985 Nov; 16(9):599–604.
23. Behrens F. General theory and principles of external fixation. *Clin Orthop Relat Res.* 1989 Apr; (241):15–23.
24. Paley D, Catagni MA, Argnani F, Villa A, Bijnedetti GB, Cattaneo R. Ilizarov treatment of tibial nonunions with bone loss. *Clin Orthop Relat Res.* 1989 Apr; (241):146–65.
25. Martínez AA, Herrera A, Pérez JM, Cuenca J, Martínez J. Treatment of humeral shaft nonunion by external fixation: a valuable option. *J Orthop Sci.* 2001; 6(3):238–41.

Conflict of Interest: Nil

Source of Support: None

Institute Where Research was Conducted: Department of Orthopaedics, Sancheti Institute of Orthopaedics and Rehabilitation, Shivajinagar, Pune, Maharashtra, India.
University Affiliation: MUHS, Nashik, Maharashtra, India.

Year of Acceptance of Thesis: 2022

How to Cite this Article: Patel J, Joshi R, Sanghavi S, Dugad M, Sonawane D, Shyam A, Sancheti P.

Hypothesis: Improved Patient Compliance and Functional Outcomes with LRS in Treating Infected Femoral Non-Unions | *Journal of Medical Thesis* | 2025 January-June; 11(1): 17-20.